

Table 10.3. Mean catch-per-unit-effort and (standard error) for fishes collected by small hoop netting in Pool 13 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.3). See methods for definitions of catch-per-unit-effort and standard error

Common name	ALL	Main channel border, unstructured	Main channel border, wing dam	Side channel border
Gizzard shad	0.12			0.29
	(0.09)			(0.22)
Common carp	0.70	0.99	0.09	0.29
	(0.51)	(0.86)	(0.09)	(0.18)
Silver chub	0.32	0.37	0.09	0.26
	(0.17)	(0.24)	(0.09)	(0.23)
Smallmouth buffalo	0.05	0.06	0.17	0.04
	(0.04)	(0.06)	(0.17)	(0.04)
Shorthead redhorse	0.00		0.08	
	(0.00)		(0.08)	
Channel catfish	36.40	29.81	56.50	45.60
	(12.79)	(17.39)	(34.58)	(19.11)
Flathead catfish	0.00		0.08	
	(0.00)		(0.08)	
White bass	0.04	0.06	0.08	
	(0.04)	(0.06)	(0.08)	
Rock bass	0.01			0.04
	(0.01)			(0.04)
Pumpkinseed	0.12	0.12		0.11
	(0.08)	(0.12)		(0.11)
Orangespotted sunfish	0.01			0.04
	(0.01)			(0.04)
Bluegill	0.58	0.77	0.33	0.33
	(0.32)	(0.54)	(0.17)	(0.19)
White crappie	0.00		0.08	
	(0.00)		(0.08)	
Black crappie	0.01			0.04
	(0.01)			(0.04)

Table 10.3 continued.

Common name	ALL	Main channel border, unstructured	Main channel border, wing dam	Side channel border
Freshwater drum	0.12	0.06	1.02	0.18
	(0.05)	(0.06)	(0.30)	(0.08)