

Table 10.5. Mean catch-per-unit-effort and (standard error) for fishes collected by small hoop netting in the Open River Reach of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.5). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Main channel border, unstructured	Main channel border, wing dam	Side channel border
Shortnose gar	0.01			0.10
	(0.01)			(0.07)
Common carp	0.77	0.82	0.44	0.45
	(0.44)	(0.50)	(0.13)	(0.19)
Smallmouth buffalo	1.01	1.14	0.58	0.10
	(0.55)	(0.63)	(0.35)	(0.05)
Black buffalo	0.75	0.81	0.15	0.31
	(0.35)	(0.40)	(0.09)	(0.12)
Blue catfish	0.18	0.20		0.10
	(0.12)	(0.14)		(0.07)
Channel catfish	5.15	3.27	0.22	19.38
	(1.62)	(1.54)	(0.10)	(7.56)
Flathead catfish	0.25	0.26		0.19
	(0.15)	(0.17)		(0.11)
Bluegill	0.01			0.06
	(0.01)			(0.04)
Longear sunfish	0.00			0.03
	(0.00)			(0.03)
Freshwater drum	0.00		0.07	0.03
	(0.00)		(0.07)	(0.03)