

Table 11.2. Mean catch-per-unit-effort and (standard error) for fishes collected by large hoop netting in Pool 8 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.2). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Main channel border, unstructured	Side channel border
Common carp	3.04		4.88
	(2.73)		(4.38)
Smallmouth buffalo	0.77	0.06	1.20
	(0.32)	(0.06)	(0.52)
Shorthead redhorse	0.19	0.19	0.18
	(0.08)	(0.09)	(0.12)
Channel catfish	1.74	2.17	1.47
	(0.58)	(1.11)	(0.64)
Flathead catfish	0.35	0.39	0.32
	(0.14)	(0.13)	(0.21)
Northern pike	0.04		0.06
	(0.04)		(0.06)
Rock bass	0.07	0.07	0.07
	(0.05)	(0.07)	(0.07)
Bluegill	3.94	1.68	5.31
	(2.02)	(1.46)	(3.13)
Largemouth bass	0.06	0.07	0.06
	(0.05)	(0.07)	(0.06)
Black crappie	1.89	0.13	2.95
	(1.36)	(0.09)	(2.19)
Yellow perch	0.03	0.07	
	(0.03)	(0.07)	
Freshwater drum	0.06	0.07	0.06
	(0.05)	(0.07)	(0.06)