

Table 11.3. Mean catch-per-unit-effort and (standard error) for fishes collected by large hoop netting in Pool 13 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.3). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Main channel border, unstructured	Main channel border, wing dam	Side channel border
Shortnose gar	0.01			0.04
	(0.01)			(0.04)
Mooneye	0.01			0.03
	(0.01)			(0.03)
Gizzard shad	0.01			0.04
	(0.01)			(0.04)
Common carp	0.08	0.06	0.17	0.11
	(0.06)	(0.06)	(0.17)	(0.11)
River carpsucker	0.05	0.06		0.03
	(0.04)	(0.06)		(0.03)
Smallmouth buffalo	0.96	1.74	-42.15	0.82
	(0.66)	(0.84)	(43.58)	(0.48)
Shorthead redhorse	0.15	0.18	0.17	0.11
	(0.08)	(0.13)	(0.17)	(0.06)
Channel catfish	0.21		0.41	0.50
	(0.11)		(0.41)	(0.28)
Flathead catfish	0.15	0.19	0.17	0.11
	(0.08)	(0.13)	(0.17)	(0.06)
White bass	0.30	0.43	0.25	0.11
	(0.12)	(0.19)	(0.25)	(0.08)
Bluegill	0.02		-2.22	0.11
	(0.03)		(2.22)	(0.06)
White crappie	0.01			0.04
	(0.01)			(0.04)
Black crappie	0.01			0.04
	(0.01)			(0.04)
Sauger	0.02			0.04
	(0.02)			(0.04)

Table 11.3 continued.

Common name	ALL	Main channel border, unstructured	Main channel border, wing dam	Side channel border
Freshwater drum	0.15	0.18	0.33	0.11
	(0.11)	(0.18)	(0.16)	(0.08)