

*Table 14.5 Mean catch-per-unit-effort and (standard error) for fishes collected by day electrofishing in the Open River Reach of the Mississippi River using fixed-site sampling in 2007. See methods for definitions of catch-per-unit-effort and standard error.*

<b>Common name</b>	<b>Main channel border, unstructured</b>	<b>Tributary mouth</b>
<b>Spotted gar</b>	0.00	0.50
	(0.00)	(0.50)
<b>Shortnose gar</b>	0.50	0.00
	(0.50)	(0.00)
<b>Goldeye</b>	1.50	0.00
	(0.50)	(0.00)
<b>Skipjack herring</b>	0.50	0.00
	(0.50)	(0.00)
<b>Gizzard shad</b>	19.50	21.50
	(14.50)	(9.06)
<b>Common carp</b>	0.50	4.00
	(0.50)	(1.68)
<b>Silver carp</b>	0.00	39.50
	(0.00)	(39.50)
<b>Emerald shiner</b>	11.00	1.50
	(11.00)	(1.19)
<b>River carpsucker</b>	0.00	1.50
	(0.00)	(1.19)
<b>Smallmouth buffalo</b>	0.00	2.25
	(0.00)	(0.95)
<b>Bigmouth buffalo</b>	0.00	2.50
	(0.00)	(1.32)
<b>Black buffalo</b>	0.50	0.00
	(0.50)	(0.00)
<b>Shorthead redhorse</b>	0.00	0.25
	(0.00)	(0.25)
<b>Channel catfish</b>	0.00	0.50
	(0.00)	(0.29)
<b>Flathead catfish</b>	0.50	0.50
	(0.50)	(0.29)
<b>Brook silverside</b>	0.00	0.25
	(0.00)	(0.25)

*Table 14.5 continued.*

<b>Common name</b>	<b>Main channel border, unstructured</b>	<b>Tributary mouth</b>
<b>White bass</b>	0.00	0.50
	(0.00)	(0.29)
<b>Green sunfish</b>	0.00	0.50
	(0.00)	(0.29)
<b>Orangespotted sunfish</b>	0.00	0.25
	(0.00)	(0.25)
<b>Bluegill</b>	0.00	0.25
	(0.00)	(0.25)
<b>Largemouth bass</b>	0.00	0.75
	(0.00)	(0.48)
<b>White crappie</b>	0.00	0.50
	(0.00)	(0.29)
<b>Sauger</b>	0.00	0.75
	(0.00)	(0.48)
<b>Freshwater drum</b>	0.50	0.00
	(0.50)	(0.00)