

*Table 16.6. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in La Grange Pool of the Illinois River using fixed-site sampling in 2007. See text for definitions of catch-per-unit-effort and standard error.*

<b>Common name</b>	<b>Tailwater</b>
<b>Shortnose gar</b>	3.85
	(2.24)
<b>Skipjack herring</b>	0.25
	(0.25)
<b>Gizzard shad</b>	16.05
	(6.50)
<b>Threadfin shad</b>	0.51
	(0.51)
<b>Common carp</b>	14.81
	(7.40)
<b>River carpsucker</b>	1.29
	(0.66)
<b>Highfin carpsucker</b>	0.26
	(0.26)
<b>Smallmouth buffalo</b>	4.65
	(2.65)
<b>Silver redhorse</b>	0.77
	(0.50)
<b>Golden redhorse</b>	0.26
	(0.26)
<b>Shorthead redhorse</b>	3.12
	(2.45)
<b>Channel catfish</b>	1.04
	(0.74)
<b>White perch</b>	3.37
	(2.06)
<b>White bass</b>	63.51
	(46.59)
<b>Yellow bass</b>	6.44
	(4.03)
<b>White perch x yellow bass hybrid</b>	3.61
	(2.35)
<b>Green sunfish</b>	1.26
	(1.26)

*Table 16.6 continued.*

<b>Common name</b>	<b>Tailwater</b>
<b>Bluegill</b>	33.73
	(11.83)
<b>White crappie</b>	0.78
	(0.78)
<b>Black crappie</b>	2.85
	(1.31)
<b>Sauger</b>	0.51
	(0.30)
<b>Freshwater drum</b>	8.32
	(4.57)