

**Table 17.5. Mean catch-per-unit-effort and (standard error) for fishes collected by mini fyke netting in the Open River Reach of the Mississippi River using fixed-site sampling in 2007. See methods for definitions of catch-per-unit-effort and standard error.**

<b>Common name</b>	<b>Main channel border, unstructured</b>	<b>Tributary mouth</b>
<b>Shortnose gar</b>	1.04	0.26
	(1.04)	(0.26)
<b>Gizzard shad</b>	1.56	0.00
	(1.56)	(0.00)
<b>Red shiner</b>	2.02	0.00
	(0.06)	(0.00)
<b>Common carp</b>	0.00	0.27
	(0.00)	(0.27)
<b>Silver carp</b>	0.00	1.99
	(0.00)	(1.64)
<b>Emerald shiner</b>	47.03	0.74
	(31.40)	(0.74)
<b>River shiner</b>	1.56	0.00
	(1.56)	(0.00)
<b>Weed shiner</b>	0.52	0.00
	(0.52)	(0.00)
<b>Channel shiner</b>	2.05	0.81
	(1.07)	(0.51)
<b>Pugnose minnow</b>	0.52	0.53
	(0.52)	(0.53)
<b>Bullhead minnow</b>	1.01	0.00
	(0.03)	(0.00)
<b>Black buffalo</b>	0.52	0.00
	(0.52)	(0.00)
<b>Blue catfish</b>	0.52	0.00
	(0.52)	(0.00)
<b>Channel catfish</b>	1.01	0.27
	(0.03)	(0.27)
<b>Flathead catfish</b>	0.52	0.27
	(0.52)	(0.27)
<b>White bass</b>	0.00	0.54
	(0.00)	(0.31)

*Table 17.5 continued.*

<b>Common name</b>	<b>Main channel border, unstructured</b>	<b>Tributary mouth</b>
<b>Orangespotted sunfish</b>	0.00	7.13
	(0.00)	(6.78)
<b>Bluegill</b>	2.60	21.21
	(2.60)	(11.17)
<b>White crappie</b>	0.00	1.81
	(0.00)	(0.88)
<b>Black crappie</b>	0.00	0.52
	(0.00)	(0.30)
<b>Sauger</b>	0.52	0.00
	(0.52)	(0.00)
<b>Freshwater drum</b>	1.56	0.26
	(1.56)	(0.26)