

*Table 6.1. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in Pool 4 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.1). See methods for definitions of catch-per-unit-effort and standard error.*

<b>Common name</b>	<b>ALL</b>	<b>Backwater, contiguous, shoreline</b>
<b>Shortnose gar</b>	1.14	1.14
	(1.14)	(1.14)
<b>Bowfin</b>	3.48	3.48
	(1.50)	(1.50)
<b>Gizzard shad</b>	10.20	10.20
	(5.70)	(5.71)
<b>Common carp</b>	1.65	1.65
	(0.83)	(0.84)
<b>Golden shiner</b>	0.09	0.09
	(0.09)	(0.09)
<b>River carpsucker</b>	0.57	0.57
	(0.57)	(0.57)
<b>Quillback</b>	0.08	0.08
	(0.08)	(0.08)
<b>Spotted sucker</b>	0.50	0.50
	(0.29)	(0.29)
<b>Silver redhorse</b>	3.36	3.36
	(1.83)	(1.83)
<b>Golden redhorse</b>	0.09	0.09
	(0.09)	(0.09)
<b>Shorthead redhorse</b>	1.64	1.64
	(1.08)	(1.08)
<b>Black bullhead</b>	0.95	0.95
	(0.94)	(0.95)
<b>Yellow bullhead</b>	0.37	0.37
	(0.29)	(0.29)
<b>Channel catfish</b>	0.35	0.35
	(0.20)	(0.20)
<b>Northern pike</b>	1.02	1.02
	(0.28)	(0.28)
<b>White bass</b>	0.25	0.25

*Table 6.1 continued.*

<b>Common name</b>	<b>ALL</b>	<b>Backwater, contiguous, shoreline</b>
	(0.24)	(0.25)
<b>Rock bass</b>	1.88	1.88
	(0.90)	(0.90)
<b>Pumpkinseed</b>	6.40	6.40
	(2.82)	(2.82)
<b>Bluegill</b>	49.73	49.73
	(26.19)	(26.23)
<b>Green sunfish x bluegill hybrid</b>	0.08	0.08
	(0.08)	(0.08)
<b>Largemouth bass</b>	0.58	0.58
	(0.37)	(0.37)
<b>Black crappie</b>	7.30	7.30
	(2.64)	(2.65)
<b>Yellow perch</b>	7.69	7.69
	(3.44)	(3.44)
<b>Sauger</b>	0.08	0.08
	(0.08)	(0.08)
<b>Walleye</b>	0.17	0.17
	(0.12)	(0.12)
<b>Freshwater drum</b>	1.66	1.66
	(1.04)	(1.04)