

Table 6.2. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in Pool 8 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.2). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
Longnose gar	0.19	0.22	
	(0.12)	(0.14)	
Shortnose gar	0.59	0.64	0.25
	(0.32)	(0.36)	(0.25)
Bowfin	1.78	1.99	0.36
	(0.50)	(0.57)	(0.25)
Gizzard shad	10.72	5.81	44.48
	(4.76)	(4.18)	(24.31)
Common carp	1.10	0.99	1.84
	(0.35)	(0.37)	(1.15)
Golden shiner	0.15	0.16	0.12
	(0.08)	(0.09)	(0.12)
Spotted sucker	0.26	0.25	0.35
	(0.13)	(0.14)	(0.17)
Silver redhorse	1.36	1.51	0.34
	(0.51)	(0.58)	(0.24)
Shorthead redhorse	0.82	0.52	2.92
	(0.30)	(0.23)	(1.82)
Yellow bullhead	0.14	0.09	0.53
	(0.10)	(0.09)	(0.53)
Channel catfish	0.33	0.24	0.96
	(0.17)	(0.13)	(0.96)
Flathead catfish	0.26	0.28	0.12
	(0.11)	(0.12)	(0.12)
Northern pike	1.50	1.67	0.34
	(0.43)	(0.49)	(0.24)
White bass	0.21	0.24	
	(0.08)	(0.09)	
Rock bass	0.50	0.48	0.67
	(0.20)	(0.22)	(0.43)
Pumpkinseed	1.27	1.35	0.70

Table 6.2 continued.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
	(0.57)	(0.65)	(0.57)
Warmouth	0.03	0.04	
	(0.03)	(0.04)	
Bluegill	92.44	96.29	66.01
	(23.28)	(26.58)	(21.01)
Pumpkinseed x bluegill hybrid	0.04	0.04	
	(0.04)	(0.04)	
Largemouth bass	2.23	1.02	10.58
	(0.72)	(0.39)	(5.03)
White crappie	0.14	0.16	
	(0.14)	(0.16)	
Black crappie	10.07	10.41	7.80
	(2.03)	(2.29)	(3.18)
Yellow perch	3.02	3.00	3.18
	(0.83)	(0.94)	(1.13)
Walleye	0.03	0.04	
	(0.03)	(0.04)	
Freshwater drum	1.00	0.86	1.95
	(0.31)	(0.33)	(0.93)