

Table 6.4. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in Pool 26 of the Mississippi River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.4). See methods for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
Spotted gar	0.14	0.15	0.11
	(0.12)	(0.15)	(0.11)
Shortnose gar	9.44	11.32	1.30
	(6.94)	(8.59)	(0.49)
Gizzard shad	14.26	17.38	0.73
	(11.05)	(13.67)	(0.29)
Common carp	0.05		0.26
	(0.03)		(0.17)
Silver carp	0.13	0.13	0.15
	(0.11)	(0.13)	(0.15)
Bighead carp	0.21	0.26	
	(0.21)	(0.26)	
River carpsucker	0.23	0.25	0.13
	(0.14)	(0.17)	(0.13)
Smallmouth buffalo	0.22	0.27	
	(0.15)	(0.18)	
Channel catfish	0.27	0.25	0.37
	(0.21)	(0.25)	(0.18)
Flathead catfish	0.03		0.15
	(0.03)		(0.15)
White perch	0.02		0.13
	(0.02)		(0.13)
White bass	4.01	2.93	8.73
	(1.45)	(1.57)	(3.81)
Yellow bass	0.13	0.13	0.13
	(0.10)	(0.13)	(0.13)
Orangespotted sunfish	0.43	0.50	0.11
	(0.31)	(0.38)	(0.11)
Bluegill	8.01	9.36	2.15
	(3.32)	(4.10)	(1.26)
White crappie	0.30	0.37	

Table 6.4 continued.

Common name	ALL	Backwater, contiguous, shoreline	Impounded, shoreline
	(0.21)	(0.26)	
Black crappie	2.11	2.39	0.93
	(1.09)	(1.34)	(0.45)
Freshwater drum	0.17	0.13	0.38
	(0.11)	(0.13)	(0.27)