

Table 6.6. Mean catch-per-unit-effort and (standard error) for fishes collected by fyke netting in La Grange Pool of the Illinois River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.6). See text for definitions of catch-per-unit-effort and standard error.

Common name	ALL	Backwater, contiguous, shoreline
Spotted gar	0.54	0.54
	(0.19)	(0.19)
Shortnose gar	7.83	7.83
	(2.02)	(2.03)
Bowfin	0.83	0.83
	(0.31)	(0.31)
Gizzard shad	7.86	7.86
	(3.83)	(3.84)
Threadfin shad	0.05	0.05
	(0.05)	(0.05)
Grass carp	0.41	0.41
	(0.31)	(0.31)
Common carp	3.31	3.31
	(1.35)	(1.35)
Silver carp	0.10	0.10
	(0.10)	(0.10)
Bighead carp	0.35	0.35
	(0.20)	(0.20)
River carpsucker	1.50	1.50
	(0.53)	(0.53)
Quillback	0.10	0.10
	(0.09)	(0.10)
Smallmouth buffalo	0.85	0.85
	(0.28)	(0.28)
Bigmouth buffalo	0.21	0.21
	(0.09)	(0.09)
Shorthead redhorse	0.25	0.25
	(0.14)	(0.14)
Black bullhead	0.29	0.29
	(0.17)	(0.17)
Brown bullhead	0.05	0.05

Table 6.6 continued.

Common name	ALL	Backwater, contiguous, shoreline
	(0.05)	(0.05)
Channel catfish	0.34	0.34
	(0.13)	(0.13)
White perch	0.14	0.14
	(0.10)	(0.10)
White bass	11.07	11.07
	(3.03)	(3.04)
Yellow bass	5.67	5.67
	(1.67)	(1.68)
White perch x yellow bass hybrid	0.86	0.86
	(0.55)	(0.55)
Warmouth	0.05	0.05
	(0.05)	(0.05)
Orangespotted sunfish	0.19	0.19
	(0.13)	(0.13)
Bluegill	5.32	5.32
	(2.36)	(2.36)
Largemouth bass	0.34	0.34
	(0.20)	(0.20)
White crappie	2.92	2.92
	(1.50)	(1.50)
Black crappie	2.10	2.10
	(1.14)	(1.14)
Freshwater drum	1.05	1.05
	(0.27)	(0.27)