

**Table 8.6. Mean catch-per-unit-effort and (standard error) for fishes collected by mini fyke netting in La Grange Pool of the Illinois River using stratified random sampling in 2007. The statistics under ALL pertain to unbiased means over ALL strata sampled by this gear (as indicated by nonmissing entries below and by Table 2.6). See text for definitions of catch-per-unit-effort and standard error.**

<b>Common name</b>	<b>ALL</b>	<b>Backwater, contiguous, shoreline</b>	<b>Main channel border, unstructured</b>	<b>Side channel border</b>
<b>Spotted gar</b>	0.03 (0.02)	0.10 (0.07)		
<b>Shortnose gar</b>	0.39 (0.15)	0.10 (0.07)	0.50 (0.22)	0.33 (0.18)
<b>Gizzard shad</b>	6.03 (2.53)	14.74 (8.90)	2.84 (1.55)	5.60 (2.97)
<b>Grass carp</b>	0.53 (0.41)	1.68 (1.58)	0.06 (0.06)	1.21 (0.71)
<b>Red shiner</b>	2.68 (2.13)		0.69 (0.34)	48.19 (46.94)
<b>Common carp</b>	2.71 (1.24)	4.31 (3.59)	1.21 (0.74)	16.72 (14.22)
<b>Silver carp</b>	70.27 (54.98)	213.61 (212.42)	17.31 (8.67)	68.97 (46.72)
<b>Bighead carp</b>	0.01 (0.01)	0.05 (0.05)		
<b>Silver chub</b>	0.00 (0.00)			0.09 (0.09)
<b>Golden shiner</b>	0.23 (0.12)	0.36 (0.36)	0.19 (0.10)	0.16 (0.11)
<b>Emerald shiner</b>	15.89 (4.96)	4.17 (2.07)	17.25 (6.49)	61.48 (43.81)
<b>River shiner</b>	0.55 (0.55)	2.14 (2.14)		
<b>Spottail shiner</b>	0.09 (0.09)		0.13 (0.13)	
<b>Silverband shiner</b>	1.52 (0.94)	0.74 (0.59)	1.78 (1.33)	1.99 (1.34)
<b>Sand shiner</b>	0.09 (0.09)		0.13 (0.13)	

*Table 8.6 continued.*

<b>Common name</b>	<b>ALL</b>	<b>Backwater, contiguous, shoreline</b>	<b>Main channel border, unstructured</b>	<b>Side channel border</b>
<b>Bullhead minnow</b>	3.33	1.45	4.11	2.05
	(1.39)	(0.60)	(1.98)	(0.72)
<b>Smallmouth buffalo</b>	0.07	0.25		
	(0.05)	(0.21)		
<b>Unidentified Catostomidae (suckers)</b>	0.01	0.05		
	(0.01)	(0.05)		
<b>Black bullhead</b>	0.02	0.05		0.08
	(0.01)	(0.05)		(0.08)
<b>Yellow bullhead</b>	0.01	0.05		
	(0.01)	(0.05)		
<b>Channel catfish</b>	2.52	0.31	3.43	1.09
	(0.62)	(0.15)	(0.89)	(0.46)
<b>Freckled madtom</b>	0.04		0.06	
	(0.04)		(0.06)	
<b>Flathead catfish</b>	0.09		0.12	0.08
	(0.06)		(0.08)	(0.08)
<b>Blackstripe topminnow</b>	0.02	0.05		0.24
	(0.01)	(0.05)		(0.17)
<b>Western mosquitofish</b>	17.82	53.15	5.53	5.81
	(13.88)	(52.49)	(4.67)	(4.61)
<b>Brook silverside</b>	0.10	0.05	0.13	
	(0.09)	(0.05)	(0.13)	
<b>White bass</b>	0.49	0.30	0.57	0.33
	(0.18)	(0.13)	(0.26)	(0.25)
<b>Yellow bass</b>	0.09	0.15	0.06	0.26
	(0.05)	(0.11)	(0.06)	(0.26)
<b>Green sunfish</b>	0.34	0.29	0.38	
	(0.12)	(0.25)	(0.15)	
<b>Warmouth</b>	0.03	0.10		
	(0.03)	(0.10)		

*Table 8.6 continued.*

<b>Common name</b>	<b>ALL</b>	<b>Backwater, contiguous, shoreline</b>	<b>Main channel border, unstructured</b>	<b>Side channel border</b>
<b>Orangespotted sunfish</b>	6.13	3.40	7.35	2.90
	(3.03)	(1.36)	(4.32)	(1.40)
<b>Bluegill</b>	3.57	7.59	2.21	1.50
	(1.30)	(4.18)	(1.05)	(0.88)
<b>White crappie</b>	0.30	0.46	0.25	0.16
	(0.11)	(0.20)	(0.15)	(0.11)
<b>Black crappie</b>	0.01			0.16
	(0.01)			(0.16)
<b>Unidentified Centrarchidae (sunfishes)</b>	0.04	0.15		
	(0.04)	(0.15)		
<b>Mud darter</b>	0.09	0.15	0.06	0.08
	(0.05)	(0.08)	(0.06)	(0.08)
<b>Sauger</b>	0.05		0.07	
	(0.05)		(0.07)	
<b>Freshwater drum</b>	0.78	0.20	1.02	0.34
	(0.31)	(0.12)	(0.44)	(0.23)